

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

March 15, 2019

Shyam Chari President Agro Logistic Systems, Inc. P.O. Box 5799 Diamond Bar, CA 91765

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Acceptable

Addition to the First Aid Statements

Product Name: Debug Turbo

EPA Registration Number: 70310-5 Application Date: 02/27/2019 OPP Decision Number: 548849

Dear Mr. Chari:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false

Page 2 of 2 EPA Reg. No. 70310-5 OPP Decision No. 548849

or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Alex Horansky by phone at (703) 347-0128 or via email at Horansky.alex@epa.gov.

Sincerely,

Andrew Bryceland, Team Leader Biochemical Pesticides Branch Biopesticides and Pollution Prevention Division (7511P) Office of Pesticide Programs

Enclosure

ACCEPTED

03/15/2019

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 70040 F

70310-5

Master Label Debug®Turbo

Page 1 of 6

Sub-label A: For Agriculture, Horticulture & Greenhouses, Golf Courses, Turf, & Ornamentals

Sub-label B: For Home Gardens

Emulsifiable Concentrate

AGRICULTURAL, HORTICULTURAL, GREENHOUSE, AND OUTDOOR DOMESTIC TERRESTRIAL PESTICIDE

ACTIVE INGREDIENT:

Fats and Glyceridic

Oils Margosa 65.80% Azadirachtin 0.70% OTHER INGREDIENTS: 33.50%

TOTAL: 100.00%

4.955 LBS (2247 GRAMS) ACTIVE INGREDIENT PER GALLON OF PRODUCT

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA REGISTRATION No. 70310-5 EPA ESTABLISHMENT No. 070310-IND-002 OR 92815-IND-001

Manufactured for:
Agro Logistic Systems, Inc.
P.O. Box 5799
Diamond Bar, CA 91765
U.S.A.

TEL: (714) 990-9220 e-mail: info@agrologistic.com Website: www.agrologistic.com

Net contents: [4 Fl. Oz.]/ [8 Fl. Oz]/ [32 Fl. Oz]/ [1 Gal]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS Harmful if absorbed through skin or if inhaled. Causes moderate eye irritation. Avoid breathing spray. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators, Mixers and other handlers must wear -Coveralls

- Socks, shoes, and
- -Chemical resistant gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables use detergent and hot water. Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This product is toxic to fish and aquatic invertebrates. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Batch No:

Manufacturing Date:

Expiry Date:

PRODUCT OF INDIA

MANUFACTURED FOR Agro Logistic Systems, Inc. P.O. Box 5799, Diamond Bar, CA 91765, U.S.A.

> TEL: (714) 990-9220 Website: www.agrologistic.com

EPA REGISTRATION No. 70310-5 EPA ESTABLISHMENT No.070310-IND-002 92815-IND-001

Sub-label A: Agriculture, Horticulture, Greenhouse, **Golf Courses, Turf, Parks & Ornamentals**

Debug®Turbo

Emulsifiable Concentrate Antifeedant, Insect Repellent, Insecticide, Miticide, Fungicide & Nematicide



FOR ORGANIC PRODUCTION

ACTIVE INGREDIENT:

Fats and Glyceridic

Oils Margosa

65.80% 0.70%

Azadirachtin OTHER INGREDIENTS:

33.50%

TOTAL:

100.00%

4.955 LBS (2247 GRAMS) ACTIVE INGREDIENT PER GALLON OF PRODUCT

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If on skin or clothing	- Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes - Call a poison control center or doctor for treatment advice.
If inhaled	 Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.
If In Eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. call a poison control center or doctor for treatment advice.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Monday through Friday, 8:00AM to 12 Noon Pacific Time. (NPIC Web Site: www.npic.orst.edu). For Medical Emergencies call your poison control center at 1-800-222-1222.	

Net contents: [4 Fl. Oz]/[8 Fl. Oz]/[32 Fl. Oz]/[1 Gal]

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Refer to the attached booklet for the listing of crops and pests treated, dilution directions and application rates.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personnel protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is:

- Coveralls
- Chemical resistant gloves, such as barrier laminates or butyl rubber or nitrile rubber or PVC.
- Shoes and socks

NON- AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used on agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. PESTICIDE STORAGE: Keep in original packaging in cool and dry place (between 60 degrees F and 80 degrees F) away from sunlight and foodstuffs. Do not refrigerate. For Commercial Agricultural Use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal

CONTAINER HANDLING: Nonrefillable Container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture or dispose of in a sanitary landfill, or by incineration. Do not burn unless allowed by state and local ordinances.

DEBUG TURBO PLEASE READ THE FOLLOWING INFORMATION CAREFULLY BEFORE USING DEBUG TURBO

TANK MIXING:

'DEBUG TURBO is compatible with commonly used insecticides, fungicides and foliar fertilizers. Do not use Debug Turbo with Bordeaux mixture, lime sulfur or other highly alkaline materials.

It is advisable to pre-test the tank mix combination proposed on a few plants to observe phytotoxicity, if any. The climatic conditions and cultural practices may be different in the actual field than experimental conditions. This product is damaged by oxidants. Do not mix acids and bleaches with DEBUG TURBO.

MIXING DIRECTIONS:

- 1. Determine the acreage to be treated with DEBUG TURBO.
- 2. Fill tank with appropriate amount of water.
- 3. Add the pre-measured amount of DEBUG TURBO.
- Keep tank agitated with an agitation device. Proper agitation is required, especially when mixing DEBUG TURBO with other pesticides.

SPRAYING DIRECTIONS:

Spray DEBUG TURBO using a boom system, propeller turbine system (indicated for orchards); pressure hose with gun sprayers, powered back sprayers and manual hand sprayers. Use a minimum boom pressure of 160 psi and maximum tractor speed of 2.0 miles per hour. Use horizontal nozzles at 10- inch spacing and two vertical drops, one at each end. Each vertical drop must have two nozzles, one at 90° angle and the other at 45° angle. The water used for mixing should have a pH at about 5.5 for DEBUG TURBO to be most effective. If necessary, use a buffer to bring the pH to the desired level.

- DURING APPLICATION, MAKE SURE THAT THE SPRAY COVERS THE ENTIRE PLANT INCLUDING THE UNDERSIDE OF LEAVES.
- AERIAL APPLICATIONS: DEBUG TURBO may be aerially applied using suitable equipment such as fixed-wing aircraft or helicopters

CHEMIGATION DIRECTIONS

GENERAL CHEMIGATION REQUIREMENTS:

- A. Apply this product only through sprinkler, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State
 Extension Service specialists, equipment manufacturers or other experts.
 5.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPRINKLER CHEMIGATION REQUIREMENTS:

 The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops". The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected" Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- 6. Agitation is generally recommended in the pesticide supply tank.
- 7. Apply this product continuously for the duration of the water application.
- A supply tank is recommended for this product. If using a supply tank, dilute this product at the rate of 1.25 gallons per 500 gallons of water.
- Apply between 1200 to 1500 gallons of water per acre to achieve efficacy.

DRIP (TRICKLE) CHEMIGATION REQUIREMENTS:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Agitation is generally recommended in the pesticide supply tank.
- 8. Apply this product continuously for the duration of the water application.
- A supply tank is recommended for this product. If using a supply tank, dilute this product at the rate of 1.25 gallons per 500 gallons of water.
- Apply between 1200 to 1500 gallons of water per acre to achieve efficacy.

For Golf Courses, Turf, Parks & Ornamentals NEMATODE, AND TURF INSECT CONTROL

For Organic Turf Use.Controls many insects including:
Red ants, Cutworms, Grubs, Chinch bugs, Crickets, Aphids, Whiteflies, Thrips,
Beetles, Leafhoppers, Fruit worms, Spider mites, Caterpillars, Cabbage
loopers, Armyworms, Colorado potato beetles, Diamondback moths,
Grasshoppers, Maggots, Weevils, Chafers, Spider Mites, spittlebugs,
Webworms, Rhizoctonia solani, Scletorium rolfsii, Sclerotinia sclerotiorum,
Fusarium oxysporum, Rust, and Powdery Mildew, Pythium

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

- 1. SHAKE THIS CONTAINER WELL BEFORE USE.
- 2A. FOR NEMATODE CONTROL: (Ornamentals and Turf) MIX 24 to 48 FL. OZ OF DEBUG TURBO WITH 2 TO 50 GALLONS WATER TO SPRAY ON 1000 SQUARE FEET. Using sprinklers, deliver additional amounts of water to push the material to the required depth below ground level. Apply every week or as needed. Repeat application if the nematode count is over the threshold limit.
- 2B. FOR INSECT AND WORM CONTROL (see above list):
 - MIX 1 to 2 FL. OZ. OF DEBUG TURBO IN 2 TO 3 GALLONS OF WATER (or 48 FL. OZ. to 96 FL. OZ. of Debug Turbo with 100 gallons water per acre). Ensure that pH of this mixture is around 5.5 by using a buffer. Spray the mixture on to the turf. Using sprinklers or other means, deliver additional amounts of water to push the material to ¼ inch ½ inch below ground level. For worm and insect control apply at the earliest sign of infestation. Repeat the application at 7-10 days interval if infestation persists.
- SPRAY DEBUG TURBO IN THE EARLY MORNING OR LATE AFTERNOON, OR EVENINGS.
- USE ALL OF THE DILUTED MATERIAL SOON AFTER MIXING. DO NOT SAVE MIXED MATERIAL.
- 5. USE AS A PREVENTATIVE TO KEEP INSECTS AWAY.

GENERAL:.

- DEBUG TURBO is more effective when used as a preventative.
- DEBUG TURBO is a broad-spectrum herbal insecticide.
- DEBUG TURBO affects the insects at all stages of growth.
- SHAKE THE CONTAINER WELL, BEFORE USE.
- IF GELLING OCCURS, WARM IT TO ROOM TEMPERATURE BEFORE APPLICATION.

WARRANTY:

AGRO LOGISTIC SYSTEMS, INC. warrants that material contained herein conforms to the description on the label and is reasonably useful for the purposes referred to in the directions for use. Method of application and timing, weather, cultural practices, insect problem, crop condition, and incompatibility with other chemicals not specifically recommended in the use of this product are beyond the control of the seller. It is the manufacturer's intention that the buyer assumes risk of use, storage or handling of this material not in strict accordance with directions given herein.

AGRICULTURAL, HORTICULTURAL, AND GREENHOUSE USE

CROPS AND APPLICATION SITES

Top fruits including but not limited to Citrus, Lemons, Oranges, Grapefruits, Pummalo, Apples, Plums, Peaches, Apricots, Avocados, Figs, Prunes, Nectarines, Pears. Mangoes, Cherries, Kiwifruit, Kumquats, Loquats, Quinces, Persimmons, and Palm.

Cucurbits, Watermelons, Gherkins, Squashes, Pumpkins, Gourds, Bitter melons, Chayote, Okra, Asparagus, and Melons. Bulb, Cole and Leafy vegetables including but not limited to Broccoli, Brussels sprouts, Cabbage, Cauliflower, Celery, Lettuce, Endives, Kale, Parsley, Onions, Garlic, Shallots, Leeks, Kohlrabi, Chard, Collard, Amaranth, Radicchio, Chicory, Mustard Greens, Rapeseed, Mizuna, Arugula, and Spinach. Root and Tuber vegetables including but not limited to Beets, Carrots, Ginger, Radish, Horseradish, Potatoes, Turnips, Yams, Sweet Potatoes, Ginseng, Rutabagas, Watercress, Artichokes and Turmeric.

Legumes including but not limited to Beans, Lentils, Peas, Peanuts, Sovbeans and Garbanzo Beans.

Cereals including but not limited to Barley, Corn, Millet, Rice, Rye, Sorghum, Triticale, Oats, and Wheat.

Alfalfa and other feed & forage crops. Grasses.

Fruiting vegetables including but not limited to Tomatoes,

Tomatillos, Bell Peppers, Chili Peppers, and Eggplant.

Berries including but not limited to Blackberries, Blueberries,

Currants, Strawberries, Cane berries, Gooseberries,

Raspberries, Cranberries, Loganberries, Huckleberries, and Boysenberries.

Grapes

Tropical fruits including but not limited to Banana, Cherimoyas,

Mangoes, Guavas, Papayas and Pineapple.

Cotton and oilseeds including but not limited to Canola,

Sesame, Rapeseed, Sunflower, Safflower, Olive, Peanut and

Jojoba

Herbs & Spices including but not limited to Mints, Anise, Basil, Chives, Coriander, Dill, Fennel, Marigold, Sage, and Thyme. Mushrooms, including but not limited to Agaricus, Bella,

Enoki, Maitake, Ovster, Shitake, and other specialty

mushrooms. Nuts including but not limited to Almonds, Pecans,

Pistachios, Cashews, Chestnuts, Macadamia, Hazelnuts, Brazil

nut, Water chestnuts, and Walnuts.

Coffee, Tea, Tobacco and Sugarcane.

Greenhouse Crops and ornamentals including but not limited to Chrysanthemums, Poinsettias, Roses, Lilies, Geraniums,

Daisies, Carnations, Salvias and Dahlias. All Starter Crops.

INSECT

PESTS/MITES/DISEASES/NEMATODES

Aphids

Armyworms

Fall Armyworms

Beetles, Beanleaf Beetles, Cucumber Beetles, Japanese Beetles, Mexican Beetles, Colorado Potato Beetles, Potato Flea Beetles, Corn beetles,

Borers, Budworms.

Casebearers, Caterpillars, Codling Moth

Fruit Flies

Flea Beetles

Gnats, Fungus Gnats, Grasshoppers, Grubs Leaf Hoppers, Leaf Perforators, Leafminers, Leafrollers, Loopers, Cabbage loopers, Lygus Bug Maggots, Onion Maggots, Mealybugs, Mildew, Rust and Powdery Mildew.

Mites, Two Spotted Spidermites, Pacific Spider Mites.

Moths, Diamondback Moths, Gypsy Moths, Grape Berry Moths,

Nematodes

Orange Totrix

Phylloxera, Phythium, Psylla, Psyllids Rhizoctonia

Solani

Scales, San Jose Scale

Sclerotinia Sclerotiorum

Sclerotium Rolfsii

Sharpshooters

Sucker Ants

Thrips

Weevils, Pepper Weevils, Boll weevils

Whiteflies

Worms

Wireworms

Webworms

Budworms

Alfalfa Worms

Boll Worms

Pickle Worms

Fagot Worms, Root Worms, Ear Worms, Cut

Worms

DIRECTIONS FOR CROP APPLICATIONS

STANDARD APPLICATION: ALL APPLICATION RATES ARE FOR ONE ACRE

16 Fl. Oz to 104 Fl. Oz. of Debug Turbo mixed with 100 gallons water or sufficient amount of water to ensure total foliar coverage. When applying in high population crops like berries and fruit trees, use dilution of 32 Fl. Oz. to 104 Fl. Oz. of Debug Turbo with 100 to 300 gallons of water.

Repeat application at 7 to 14 day intervals, at least once and if necessary additional

For Greenhouse crops Prepare a 0.50% - 1.00% (0.60 oz - 1.25 oz./ Gal) solution of Debug Turbo in water and completely drench the plants. Repeat every 5 to 7 days till infestation clears. For NEMATODE CONTROL use 104 oz. Debug Turbo per acre per application. If using Drip Irrigation add enough water to deliver the product to the root zone. If using Sprinklers apply the product by sprayer in the planting bed. Then use Sprinklers to deliver the product with enough water to reach the root zone. Repeat applications at the same rate above, as needed.

For Milk Rooms, and Barns apply at the rate of not more than 1 Fl. Oz. per 420 Sq. Ft. on the floors, after dissolving in sufficient quantity of water.

WE RECOMMEND USING DEBUG TURBO AS PREVENTATIVE APPLICATION

EVEN IF THERE IS NO INFESTATION

Start applying when the plants are tall enough to be susceptible to infestation. Use the following:

- For Baby Vegetables, apply 10 days after planting and repeat again in 7 to 10 days.
- For Regular Vegetables, apply when the plant has grown two leaves or is 6 inches tall. Repeat application every 10 to 15 days.
- For Fruit Trees, apply when the tree ends dormancy and continue to repeat applications every 10 to 15 days.
- For **Berries**, apply when the plant ends dormancy and continue to repeat applications every 10 to 15 days.
- For Grasses and Forage Crops like Alfalfa, apply in spring, approximately 2 weeks after dormancy period is over, and repeat every 10 to 15 days.

MAXIMUM APPLICATION RATE IS 20 GRAMS OF ACTIVE INGREDIENT (104 FL. OZ. OF DEBUG TURBO E.C.) OR LESS PER ACRE ACCORDING TO TOLERENCE EXEMPTION 40 CFR 180.1119 FOR AZADIRACHTIN. FATS, AND GLYCERIDIC OILS MARGOSA IS EXEMPT FROM THE REQUIREMENT OF A TOLERANCE AS PER 40 CFR 180.1291. THERE IS NO LIMIT ON THE NUMBER OF APPLICATIONS.

Agro Logistic Systems, Inc. P.O.Box: 5799 Diamond Bar, CA 91765, U.S.A. TEL: (714) 990-9220; FAX: (714) 990-9222

e-mail: info@agrologistic.com website: www.agrologistic.com

EPA REGISTRATION NO: 70310-5 EPA ESTABLISHMENT NO: 070310-IND-002 OR 92815-IND-001

Sub-label B: For Home Gardens Debug®Turbo

For Indoor and Outdoor Use For Organic Gardening

From the Neem Tree

Ideal for Flower Beds, Vegetable Gardens, Lawns & Fruit Trees Including indoor and outdoor plants grown in containers.

Antifeedant, Insect Repellent, Insecticide, Miticide, Fungicide & Nematicide Controls many insects including:

Red ants, Cutworms, Grubs, Chinch bugs, Crickets, Aphids, Whiteflies, Thrips, Beetles, Leafhoppers, Fruit worms, Spider mites, Caterpillars, loopers, Armyworms, Grasshoppers, Maggots, Weevils, Spider Mites, spittlebugs, Webworms, Rhizoctonia solani, Scletorium rolfsii, Sclerotinia sclerotiorum, Fusarium oxysporum, Rust, and Powdery Mildew, Pythium etc.

KEEP OUT OF REACH OF CHILDREN **CAUTION**

FIRST AID		
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice. 	
If inhaled	 Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. 	
If In Eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. call a poison control center or doctor for treatment advice. 	
HOT LINE NUMBER		

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Monday through Friday, 8:00AM to 12 Noon Pacific Time. (NPIC Web Site: www.npic.orst.edu). For Medical Emergencies call your poison control center at 1-800-222-1222.

ACTIVE INGREDIENT:

Fats and Glyceridic

Oils Margosa Azadirachtin

65.80% 0.70%

OTHER INGREDIENTS:

33.50%

TOTAL:

100.00%

4.955 LBS (2247GRAMS) ACTIVE INGREDIENT PER GALLON OF PRODUCT

Net contents: [0.16 Fl. Oz]/[1 Fl. Oz]/[4 Fl. Oz]/[8 Fl. Oz]/[16 Fl. Oz]/[24 Fl. Oz]/[32 Fl. Oz]

EPA REGISTRATION No. 70310-5 EPA ESTABLISHMENT No. 070310-IND-002 OR

92815-IND-001

Manufactured for: Agro Logistic Systems, Inc.

P.O. Box 5799

Diamond Bar, CA 91765, U.S.A.

Tel: (714) 990-9220

PLEASE REFER TO THE FRONT PANEL FOR <u>FIRST AID INSTRUCTIONS</u> PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through skin or if inhaled. Causes moderate eye irritation. Avoid breathing spray. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE) required for this product are long-sleeve shirt and long pants, socks, shoes, and chemical resistant gloves.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This product is toxic to fish and aquatic invertebrates. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

- 1. SHAKE THIS CONTAINER WELL BEFORE USE.
- 2. MIX 15 ML to 30 ML (1 to 2 TABLEPOONS) OF DEBUG TURBO WITH 24 TO 32 FL. OZ. WATER IN A SPRAY CONTAINER, SHAKE WELL.
- 3. SPRAY THE MIXTURE ON THE LEAVES AND STEMS, INCLUDING THE UNDERSIDE OF LEAVES SO AS TO COMPLETELY DRENCH THE PLANT.
- 4. SPRAY DEBUG TURBO EARLY IN THE MORNING OR LATE AFTERNOON.
- 5. USE ALL OF THE DILUTED MATERIAL SOON AFTER MIXING. DO NOT SAVE MIXED MATERIAL.
- 6. FOR BEST RESULTS, APPLY WEEKLY (OR MORE FREQUENTLY IF INFESTATION PERSISTS).
- 7. USE AS PREVENTATIVE TO KEEP INSECTS AWAY.
- 8. DO NOT WASH THE LEAVES FOR 24 HOURS AFTER SPRAYING.
- 9. FOR LAWNCARE, APPLY 30 ml (2 TABLESPOONS) TO 60 ml (4 TABLESPOONS) OR LESS OF CONCENTRATE PER 100 SQ. FT. PER APPLICATION.
- 10. IF GELLING OCCURS, WARM IT TO ROOM TEMPERATURE BEFORE APPLICATION

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in original packaging in cool and dry place away from sunlight and foodstuffs. Do not refrigerate. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY STATEMENT

AGRO LOGISTIC SYSTEMS, INC. warrants that material contained herein conforms to the description on the label and is reasonably useful for the purposes referred to in the directions for use. Method of application and timing, weather, cultural practices, insect problem, crop condition, and incompatibility with other chemicals not specifically recommended in the use of this product are beyond the control of the seller. It is the manufacturer's intention that the buyer assumes risk of use, storage, or handling of this material not in strict accordance with directions given herein.

Manufacturing Date: Expiry Date:

Product of India